arXiv:0907.3012v1 [math.DG] 17 Jul 2009

Lower dimensional volumes and the Kastler-Kalau-Walze type theorem for Manifolds with Boundary

Yong Wang

Abstract

In this paper, we define lower dimensional volumes of spin manifolds with boundary. We compute the lower dimensional volume $\mathrm{Vol}^{(2,2)}$ for 5-dimensional and 6-dimensional spin manifolds with boundary and we also get the Kastler-Kalau-Walze type theorem in this case.

MSC: 58G20; 53A30; 46L87

Keywords: Lower dimensional volumes, noncommutative residue for manifolds with boundary; gravitational action for manifolds with boundary

1 Introduction

The noncommutative residue found in [Gu] and [Wo] plays a prominent role in noncommutative geometry. In [C1], Connes used the noncommutative residue to derive a conformal 4-dimensional Polyakov action analogy. In [C2], Connes proved that the noncommutative residue on a compact manifold M coincided with the Dixmier's trace on pseudodifferential operators of order $-\mathrm{dim}M$. Several years ago, Connes made a challenging observation that the noncommutative residue of the square of the inverse of the Dirac operator was proportional to the Einstein-Hilbert action, which we call the Kastler-Kalau-Walze theorem. In [K], Kastler gave a brute-force proof of this theorem. In [KW], Kalau and Walze proved this theorem in the normal coordinates system simultaneously. In [A], Ackermann gave a note on a new proof of this theorem by means of the heat kernel expansion.

On the other hand, Fedosov et al defined a noncommutative residue on Boutet de Monvel's algebra and proved that it was a unique continuous trace in [FGLS]. In [Wa1] and [Wa2], we generalized some results in [C1] and [U] to the case of manifolds with boundary . In [Wa3], We proved a Kastler-Kalau-Walze type theorem for the Dirac operator and the signature operator for $3,4$-dimensional manifolds with boundary. Recently, Ponge defined lower dimensional volumes of Riemannian manifolds by the Wodzicki residue in [Po]. The motivation of this paper is to find a Kastler-Kalau-Walze type theorem for higher dimensional manifolds with boundary and generalize the definition of lower dimensional volumes to manifolds with boundary.

This paper is organized as follows: In Section 2, we define lower dimensional volumes of spin manifolds with boundary. In Section 3, for 6-dimensional spin manifolds with boundary and the associated Dirac operator D, we compute the lower dimensional volume $\mathrm{Vol}_6^{(2,2)}$ and get a Kastler-Kalau-Walze type theorem in this case. In Section 4, when ∂M is flat, we can define $\int_{\partial M}\mathrm{res}_{2,2}(D^{-2},D^{-2})$ and $\int_{\partial M}\mathrm{res}_{2,3}(D^{-2},D^{-2})$ (see Section 3) and get that the gravitational action for ∂M is proportional to $\int_{\partial M}\mathrm{res}_{2,2}(D^{-2},D^{-2})$ and $\int_{\partial M}\mathrm{res}_{2,3}(D^{-2},D^{-2})$, which gives two kinds of operator theoretic explanations of the gravitational action for boundary. For 5-dimensional spin manifolds with boundary and the associated Dirac operator D, we compute the lower dimensional volume $\mathrm{Vol}_5^{(2,2)}$.

2 Lower dimensional volumes of spin manifolds with boundary

In order to define lower dimensional volumes of spin manifolds with boundary, we need some basic facts and formulae about Boutet de Monvel's calculus and the definition of noncommutative residue for manifolds with boundary. We can find them in Section 2,3 in [Wa1] and Section 2.1 in [Wa3].

Let M be a n-dimensional compact oriented spin manifold with boundary ∂M. We assume that the metric g^M on M has the following form near the boundary,

$$g^M=\frac{1}{h(x_n)}g^{\partial M}+dx_n^2, \tag{2.1}$$

where $g^{\partial M}$ is the metric on ∂M. $h(x_n)\in C^\infty([0,1))=\{\widetilde{h}|_{[0,1)}|\widetilde{h}\in C^\infty((-\varepsilon,1))\}$ for some $\varepsilon>0$ and satisfies $h(x_n)>0,\ h(0)=1$ where x_n denotes the normal directional coordinate. Let $U\subset M$ be a collar neighborhood of ∂M which is diffeomorphic to $\partial M\times[0,1)$. By the definition of $C^\infty([0,1))$ and $h>0$, there exists $\widetilde{h}\in C^\infty((-\varepsilon,1))$ such that $\widetilde{h}|_{[0,1)}=h$ and $\widetilde{h}>0$ for some sufficiently small $\varepsilon>0$. Then there exists a metric $\widehat{g}$ on $\widehat{M}=M\cup_{\partial M}\partial M\times(-\varepsilon,0]$ which has the form on $U\cup_{\partial M}\partial M\times(-\varepsilon,0]$

$$\widehat{g}=\frac{1}{\widetilde{h}(x_n)}g^{\partial M}+dx_n^2, \tag{2.2}$$

such that $\widehat{g}|_M=g$. We fix a metric $\widehat{g}$ on the $\widehat{M}$ such that $\widehat{g}|_M=g$. We can get the spin structure on $\widehat{M}$ by extending the spin structure on M. Let D be the Dirac operator associated to $\widehat{g}$ on the spinors bundle $S(T\widehat{M})$. Let p_1,p_2 be nonnegative integers and $p_1+p_2\leq n$.

Definition 2.1 Lower dimensional volumes of spin manifolds with boundary are defined by $\mathrm{Vol}_n^{(p_1,p_2)}M:=\widetilde{\mathrm{Wres}}[\pi^+D^{-p_1}\circ\pi^+D^{-p_2}]$ (for the related definitions, see [Wa1], Section 2, 3).

Denote by $\sigma_l(A)$ the l-order symbol of an operator A. By (2.1.4)-(2.1.8) in [Wa3], we get

$$\widetilde{\mathrm{Wres}}[\pi^+D^{-p_1}\circ\pi^+D^{-p_2}]=\int_M\int_{|\xi|=1}\mathrm{trace}_{S(TM)}[\sigma_{-n}(D^{-p_1-p_2})]\sigma(\xi)dx+\int_{\partial M}\Phi, \tag{2.3}$$

where

$$\Phi=\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{j,k=0}^{\infty}\sum\frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!}$$
$$\times\mathrm{trace}_{S(TM)}[\partial^j_{x_n}\partial^\alpha_{\xi'}\partial^k_{\xi_n}\sigma^+_r(D^{-p_1})(x',0,\xi',\xi_n)\times\partial^\alpha_{x'}\partial^{j+1}_{\xi_n}\partial^k_{x_n}\sigma_l(D^{-p_2})(x',0,\xi',\xi_n)]d\xi_n\sigma(\xi')dx', \tag{2.4}$$

where the sum is taken over $r-k-|\alpha|+l-j-1=-n,\quad r\leq -p_1, l\leq -p_2$. Since $[\sigma_{-n}(D^{-p_1-p_2})]|_M$ has the same expression as $\sigma_{-n}(D^{-p_1-p_2})$ in the case of manifolds without boundary, so locally we can use the computations in [K], [KW], [Po] to compute the first term. The following proposition is the motivation of the definition of lower dimensional volumes of spin manifolds with boundary.

Proposition 2.2 1) *When* $p_1+p_2=n$, *then* $\mathrm{Vol}_n^{(p_1,p_2)}M=c_0\mathrm{Vol}_M$.
2) when $p_1+p_2\equiv n\bmod 1$, $\mathrm{Vol}_n^{(p_1,p_2)}M=\int_{\partial M}\Phi$.
3)

$$\mathrm{Vol}_4^{(1,1)}=-\frac{\Omega_4}{3}\int_M s\mathrm{dvol}_M;\quad \mathrm{Vol}_3^{(1,1)}=c_1\mathrm{Vol}_{\partial M} \tag{2.5}$$

where c_0, c_1 *are constants and* s *is the scalar curvature.*

Proof. 1) comes from (2.4) and (2.2) in [Po]. 2) comes from the proposition 2.3 and 3.2 in [Po]. 3) comes from Theorems 2.5 and 5.1 in [Wa3]. □

3 A Kastler-Kalau-Walze type theorem for 6-dimensional spin manifolds with boundary

In this section, We compute the lower dimensional volume $\mathrm{Vol}_6^{(2,2)}$ for 6-dimensional spin manifolds with boundary and get a Kastler-Kalau-Walze type theorem in this case.

Firstly, we recall the symbol expansion of D^{-2} in [Ka]. Recall the definition of the Dirac operator D (see [BGV], [Y]). Let ∇^L denote the Levi-civita connection about g^M. In the local coordinates $\{x_i; 1\leq i\leq n\}$ and the fixed orthonormal frame $\{\widetilde{e_1},\cdots,\widetilde{e_n}\}$, the connection matrix $(\omega_{s,t})$ is defined by

$$\nabla^L(\widetilde{e_1},\cdots,\widetilde{e_n})=(\widetilde{e_1},\cdots,\widetilde{e_n})(\omega_{s,t}). \tag{3.1}$$

$c(\widetilde{e_i})$ denotes the Clifford action. The Dirac operator

$$D=\sum_{i=1}^n c(\partial_i)(\partial_i+\delta_i);\ \delta_i=-\frac{1}{4}\sum_{s,t}\omega_{s,t}(\partial_i)c(\widetilde{e_s})c(\widetilde{e_t}). \tag{3.2}$$

Let $g^{ij} = g(dx_i, dx_j)$ and

$$\nabla^L_{\partial_i}\partial_j = \sum_k \Gamma^k_{ij}\partial_k;\ \Gamma^k = g^{ij}\Gamma^k_{ij},\ \delta^j = g^{ij}\delta_i. \tag{3.3}$$

Let the cotangent vector $\xi = \sum \xi_j dx_j$ and $\xi^j = g^{ij}\xi_i$. Then we have

Lemma 3.1([Ka])

$$\sigma_{-2}(D^{-2}) = |\xi|^{-2};\ \sigma_{-3}(D^{-2}) = -\sqrt{-1}|\xi|^{-4}\xi_k(\Gamma^k - 2\delta^k) - \sqrt{-1}|\xi|^{-6}2\xi^j\xi_\alpha\xi_\beta\partial_j g^{\alpha\beta}. \tag{3.4}$$

Since Φ is a global form on ∂M, so for any fixed point $x_0 \in \partial M$, we can choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\widetilde{U} = U \times [0, 1) \subset M$ and the metric $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$. For details, see Section 2.2.2 in [Wa3].

Now we can compute Φ (see formula (2.4) for the definition of Φ), since the sum is taken over $-r-l+k+j+|\alpha| = -5,\ \ r, l \leq -2$, then we have the following five cases:

case a) I) $r = -2,\ l = -2\ k = j = 0,\ |\alpha| = 1$

By (2.4), we get

$$\text{case a) I)} = -\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{|\alpha|=1}\mathrm{trace}[\partial^\alpha_{\xi'}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial^\alpha_{x'}\partial_{\xi_n}\sigma_{-2}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx', \tag{3.5}$$

By Lemma 2.2 in [Wa3], for $i < n$, then

$$\partial_{x_i}\sigma_{-2}(D^{-2})(x_0) = \partial_{x_i}(|\xi|^{-2})(x_0) = -\frac{\partial_{x_i}(|\xi|^2)(x_0)}{|\xi|^4} = 0, \tag{3.6}$$

so case a) I) vanishes.

case a) II) $r = -2,\ l = -2\ k = |\alpha| = 0,\ j = 1$

By (2.4), we get

$$\text{case a) II)} = -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{x_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial^2_{\xi_n}\sigma_{-2}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx', \tag{3.7}$$

By Lemma 2.2 in [Wa3], we have

$$\partial_{x_n}\sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1} = -\frac{h'(0)}{(1+\xi_n^2)^2} \tag{3.8}$$

By (3.8) and the Cauchy integral formula and (2.1.1) in [Wa3], then

$$\pi^+_{\xi_n}\partial_{x_n}\sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1} \quad = \quad -h'(0)\frac{1}{2\pi i}\mathrm{lim}_{u\to 0^-}\int_{\Gamma^+}\frac{\frac{1}{(\eta_n+i)^2(\xi_n+iu-\eta_n)}}{(\eta_n-i)^2}d\eta_n$$

$$= \quad h'(0)\frac{i\xi_n+2}{4(\xi_n-i)^2}, \tag{3.9}$$

$$\partial^2_{\xi_n}(|\xi|^{-2})(x_0)=\frac{-2+6\xi_n^2}{(1+\xi_n^2)^3}. \tag{3.10}$$

We note that

$$\begin{aligned}\int_{-\infty}^{\infty}\frac{i\xi_n+2}{(\xi_n-i)^2}\times\frac{-1+3\xi_n^2}{(1+\xi_n^2)^3}d\xi_n &= \int_{\Gamma^+}\frac{3i\xi_n^3+6\xi_n^2-i\xi_n-2}{(\xi_n-i)^5(\xi_n+i)^3}d\xi_n \\ &= \frac{2\pi i}{4!}\left[\frac{3i\xi_n^3+6\xi_n^2-i\xi_n-2}{(\xi_n+i)^3}\right]^{(4)}|_{\xi_n=i}=\frac{5\pi}{16}, \end{aligned}\tag{3.11}$$

Since $n=6$, $\mathrm{tr}_{S(TM)}[\mathrm{id}]=\dim(\wedge^*(3))=8$. So by (3.7)-(3.11), we get **case a** II)$=-\frac{5}{8}\pi h'(0)\Omega_4dx'$, where Ω_4 is the canonical volume of S^4.

case a) III) $r=-2,\ l=-2\ j=|\alpha|=0,\ k=1$

By (2.4) and an integration by parts, we get

$$\text{case a) III)}=-\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial_{\xi_n}\partial_{x_n}\sigma_{-2}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx'$$

$$=\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial^2_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial_{x_n}\sigma_{-2}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.12}$$

By Lemma 2.2 in [Wa3] , we have

$$\partial^2_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1}=\frac{-i}{(\xi_n-i)^3}. \tag{3.13}$$

By (3.8) and (3.13), we have

$$\textbf{case a) III)}=4ih'(0)\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\int_{\Gamma^+}\frac{1}{(\xi_n-i)^5(\xi_n+i)^2}d\xi_n\sigma(\xi')dx'=\frac{5}{8}\pi h'(0)\Omega_4dx'.$$

Thus the sum of **case a) II)** and **case a) III)** is zero.

case b) $r=-2,\ l=-3,\ k=j=|\alpha|=0$

By (2.4) and an integration by parts, we get

$$\text{case b)}=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial_{\xi_n}\sigma_{-3}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx'$$

$$=i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\sigma_{-3}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.14}$$

By Lemma 2.2 in [Wa3], we have

$$\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1}=\frac{i}{2(\xi_n-i)^2}. \tag{3.15}$$

In the normal coordinate, $g^{ij}(x_0)=\delta_i^j$ and $\partial_{x_j}(g^{\alpha\beta})(x_0)=0$, if $j<n$; $=h'(0)\delta^\alpha_\beta$, if $j=n$. So by Lemma A.2 in [Wa3], we have $\Gamma^n(x_0)=\frac{5}{2}h'(0)$ and $\Gamma^k(x_0)=0$ for $k<n$. By the definition of δ^k and Lemma 2.3 in [Wa3], we have $\delta^n(x_0)=0$ and $\delta^k=\frac{1}{4}h'(0)c(\widetilde{e_k})c(\widetilde{e_n})$ for $k<n$. So

$$\begin{aligned}\sigma_{-3}(D^{-2})(x_0)|_{|\xi'|=1}&=-\sqrt{-1}|\xi|^{-4}\xi_k(\Gamma^k-2\delta^k)(x_0)|_{|\xi'|=1}-\sqrt{-1}|\xi|^{-6}2\xi^j\xi_\alpha\xi_\beta\partial_jg^{\alpha\beta}(x_0)|_{|\xi'|=1}\\&=\frac{-i}{(1+\xi_n^2)^2}\left(-\frac{1}{2}h'(0)\sum_{k<n}\xi_kc(\widetilde{e_k})c(\widetilde{e_n})+\frac{5}{2}h'(0)\xi_n\right)-\frac{2ih'(0)\xi_n}{(1+\xi_n^2)^3}. \end{aligned}\tag{3.16}$$

We note that $\int_{|\xi'|=1}\xi_1\cdots\xi_{2q+1}\sigma(\xi')=0$, so the first term in (3.16) has no contribution for computing case b).

$$\begin{aligned}\textbf{case b)}\quad&=\quad ih'(0)\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}\left[\frac{1}{2(\xi_n-i)^2}\times\left(\frac{\frac{5}{2}\xi_n}{(1+\xi_n^2)^2}+\frac{2\xi_n}{(1+\xi_n^2)^3}\right)\right]d\xi_n\sigma(\xi')dx'\\&=\quad 2ih'(0)\Omega_4\int_{\Gamma^+}\frac{5\xi_n^3+9\xi_n}{(\xi_n-i)^5(\xi_n+i)^3}d\xi_ndx'\\&=\quad 2ih'(0)\Omega_4\frac{2\pi i}{4!}\left[\frac{5\xi_n^3+9\xi_n}{(\xi_n+i)^3}\right]^{(4)}|_{\xi_n=i}dx'=-\frac{15}{8}\pi h'(0)\Omega_4dx'.\end{aligned}\tag{3.17}$$

case c) $r=-3,\ l=-2,\ k=j=|\alpha|=0$

By (2.4), we get

$$\text{case c)}=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.18}$$

By the Leibniz rule, trace property and "++" and "- -" vanishing after the integration over ξ_n (for details, see [FGLS]), then

$$\begin{aligned}&\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n\\=&\int_{-\infty}^{+\infty}\mathrm{tr}[\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n-\int_{-\infty}^{+\infty}\mathrm{tr}[\pi^-_{\xi_n}\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n\\=&\int_{-\infty}^{+\infty}\mathrm{tr}[\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n-\int_{-\infty}^{+\infty}\mathrm{tr}[\pi^-_{\xi_n}\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n\\=&\int_{-\infty}^{+\infty}\mathrm{tr}[\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n-\int_{-\infty}^{+\infty}\mathrm{tr}[\sigma_{-3}(D^{-2})\times\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n\end{aligned}$$

$$
\begin{aligned}
&= \int_{-\infty}^{+\infty}\mathrm{tr}[\partial_{\xi_n}\sigma_{-2}(D^{-2})\times\sigma_{-3}(D^{-2})]d\xi_n+\int_{-\infty}^{+\infty}\mathrm{tr}[\partial_{\xi_n}\sigma_{-3}(D^{-2})\times\pi^+_{\xi_n}\sigma_{-2}(D^{-2})]d\xi_n \\
&= \int_{-\infty}^{+\infty}\mathrm{tr}[\partial_{\xi_n}\sigma_{-2}(D^{-2})\times\sigma_{-3}(D^{-2})]d\xi_n+\int_{-\infty}^{+\infty}\mathrm{tr}[\pi^+_{\xi_n}\sigma_{-2}(D^{-2})\times\partial_{\xi_n}\sigma_{-3}(D^{-2})]d\xi_n \qquad (3.19)
\end{aligned}
$$

By (3.19), we have

$$\textbf{case c)}=\textbf{case b)}-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{tr}[\partial_{\xi_n}\sigma_{-2}(D^{-2})\times\sigma_{-3}(D^{-2})]d\xi_n\sigma(\xi')dx'. \quad (3.20)$$

We note that we can not get the sum of **case b)** and **case c)** is zero by computations in (3.19). In order to compute **case c)**, we only need compute the last term in (3.20). By (3.16) and

$$\partial_{\xi_n}\sigma_{-2}(D^{-2})(x_0)|_{|\xi'|=1}=-\frac{2\xi_n}{(\xi_n^2+1)^2}, \quad (3.21)$$

we have

$$
\begin{aligned}
&-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{tr}[\partial_{\xi_n}\sigma_{-2}(D^{-2})\times\sigma_{-3}(D^{-2})]d\xi_n\sigma(\xi')dx' \\
&=8h'(0)\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{5\xi_n^4+9\xi_n^2}{(\xi_n+i)^5(\xi_n-i)^5}d\xi_n\sigma(\xi')dx'=\frac{15}{4}\pi h'(0)\Omega_4dx'. \qquad (3.22)
\end{aligned}
$$

By (3.17), (3.20) and (3.22), we have the sum of **case b)** and **case c)** is zero. Now Φ is the sum of the cases a), b) and c), so is zero. Then we get

Theorem 3.2 *Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above and D be the Dirac operator on $\widehat{M}$, then*

$$\mathrm{Vol}_6^{(2,2)}=\widetilde{\mathrm{Wres}}[(\pi^+D^{-2})^2]=-\frac{5\Omega_6}{3}\int_M s\mathrm{dvol}_M. \quad (3.23)$$

4 The gravitational action for 6-dimensional manifolds with boundary

Firstly, we recall the Einstein-Hilbert action for manifolds with boundary (see [H] or [B]),

$$I_{\mathrm{Gr}}=\frac{1}{16\pi}\int_M s\mathrm{dvol}_M+2\int_{\partial M}K\mathrm{dvol}_{\partial_M}:=I_{\mathrm{Gr,i}}+I_{\mathrm{Gr,b}}, \quad (4.1)$$

where

$$K=\sum_{1\leq i,j\leq n-1}K_{i,j}g^{i,j}_{\partial M};\quad K_{i,j}=-\Gamma^n_{i,j}, \quad (4.2)$$

and $K_{i,j}$ is the second fundamental form, or extrinsic curvature. Taking the metric in Section 2, then by Lemma A.2 in [Wa3], for $n=6$, then

$$K(x_0)==-\frac{5}{2}h'(0);\ I_{\mathrm{Gr,b}}=-5h'(0)\mathrm{Vol}_{\partial M}. \quad (4.3)$$

Let M be 6-dimensional manifolds with boundary and P, P' be two pseudodifferential operators with transmission property (see [Wa1] or [RS]) on $\widehat{M}$. Motivated by (2.4), we define locally

$$\mathrm{res}_{2,2}(P,P') := -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{x_n}\pi^+_{\xi_n}\sigma_{-2}(P)\times\partial^2_{\xi_n}\sigma_{-2}(P')]d\xi_n\sigma(\xi')dx'; \quad (4.4)$$

$$\mathrm{res}_{2,3}(P,P') := -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-2}(P)\times\partial_{\xi_n}\sigma_{-3}(P')]d\xi_n\sigma(\xi')dx'. \quad (4.5)$$

By (4.4),(4.5), so

$$\text{case a) II)} = \mathrm{res}_{2,2}(D^{-2},D^{-2});\ \text{case b)} = \mathrm{res}_{2,3}(D^{-2},D^{-2}). \quad (4.6)$$

Now, we assume ∂M is flat, then $\{dx_i = e_i\}$, $g^{\partial M}_{i,j} = \delta_{i,j}$, $\partial_{x_s}g^{\partial M}_{i,j} = 0$. So $\mathrm{res}_{2,2}(D^{-2},D^{-2})$ and $\mathrm{res}_{2,3}(D^{-2},D^{-2})$ are two global forms locally defined by the aboved oriented orthonormal basis $\{dx_i\}$. By case a) II) and case b), then we have:

Theorem 4.1 *Let M be a 6-dimensional compact spin manifold with the boundary ∂M and the metric g^M as above and D be the Dirac operator on $\widehat{M}$. Assume ∂M is flat, then*

$$\int_{\partial M}\mathrm{res}_{2,2}(D^{-2},D^{-2}) = \frac{\pi}{8}\Omega_4 I_{\mathrm{Gr,b}}; \quad (4.7)$$

$$\int_{\partial M}\mathrm{res}_{2,3}(D^{-2},D^{-2}) = \frac{3\pi}{8}\Omega_4 I_{\mathrm{Gr,b}}. \quad (4.8)$$

Nextly, for 5-dimensional spin manifolds with boundary, we compute $\mathrm{Vol}_5^{(2,2)}$. By Proposition 2.2 (2), we have

$$\widetilde{\mathrm{Wres}}[(\pi^+D^{-2})^2] = \int_{\partial M}\Phi. \quad (4.9)$$

When $n = 5$, then in (2.4), $r - k - |\alpha| + l - j - 1 = -5,\quad r, l \leq -2$, so we get $r = l = -2,\ k = |\alpha| = j = 0$, then

$$\Phi = \int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}_{S(TM)}[\sigma^+_{-2}(D^{-2})(x',0,\xi',\xi_n)\times\partial_{\xi_n}\sigma_{-2}(D^{-2})(x',0,\xi',\xi_n)]d\xi_5\sigma(\xi')dx'. \quad (4.10)$$

By (3.21) and $\pi^+_{\xi_n}\sigma_{-2}(x_0)|_{|\xi'|=1} = \frac{1}{2i(\xi_n-i)}$ and $\mathrm{tr}(id) = \dim(S(TM)) = 4$, we can get $\mathrm{Vol}_5^{(2,2)} = \frac{\pi i}{2}\Omega_3\mathrm{Vol}_{\partial M}$. By $I_{\mathrm{Gr,b}} = -4h'(0)\mathrm{Vol}_{\partial M}$, we have

Theorem 4.2 *Let M be a 5-dimensional compact spin manifold with the boundary ∂M and the metric g^M as in Section 2 and D be the Dirac operator on $\widehat{M}$, then*

$$\mathrm{Vol}_5^{(2,2)} = \widetilde{\mathrm{Wres}}[(\pi^+D^{-2})^2] = \frac{\pi i}{2}\Omega_3\mathrm{Vol}_{\partial M}, \quad (4.11)$$

$$I_{\mathrm{Gr,b}} = \frac{8ih'(0)}{\pi\Omega_3}\widetilde{\mathrm{Wres}}[(\pi^+D^{-2})^2], \tag{4.12}$$

where $\mathrm{Vol}_{\partial M}$ *denotes the canonical volume of* ∂M*.*

Remark 4.3 By Theorem 4.2, we know that $\widetilde{\mathrm{Wres}}[(\pi^+D^{-1})^2]$ is proportional to the gravitational action for boundary for 5-dimensional manifolds with boundary. But the constant depends on $h'(0)$.

Acknowledgement: The author is indebted to Professor Weiping Zhang for the suggestion to consider this problem. This work was supported by NSFC No.10801027.

References

[A] T. Ackermann, *A note on the Wodzicki residue,* J. Geom. Phys., 20, 404-406, 1996.
[B] N. H. Barth, *The fourth-order gravitational action for manifolds with boundaries,* Class. Quantum Grav. 2, 497-513, 1985.
[BGV] N. Berline, E. Getzler, M. Vergne, *Heat Kernals and Dirac Operators,* Springer-Verlag, Berlin, 1992.
[C1] A. Connes, *Quantized calculus and applications,* XIth International Congress of Mathematical Physics (paris,1994), 15-36, Internat Press, Cambridge, MA, 1995.
[C2] A. Connes. *The action functinal in noncommutative geometry,* Comm. Math. Phys., 117:673-683, 1998.
[FGLS] B. V. Fedosov, F. Golse, E. Leichtnam, and E. Schrohe. *The noncommutative residue for manifolds with boundary,* J. Funct. Anal, 142:1-31,1996.
[Gu] V.W. Guillemin, *A new proof of Weyl's formula on the asymptotic distribution of eigenvalues*, Adv. Math. 55 no.2, 131-160, 1985.
[H] S. W. Hawking, *General Relativity. An Einstein Centenary Survey,* Edited by S. W. Hawking and W. Israel, Cambridge University Press,Cambridge-New York, 1979.
[K] D. Kastler, *The Dirac operator and gravitiation,* Commun. Math. Phys, 166:633-643, 1995.
[KW] W. Kalau and M.Walze, *Gravity, non-commutative geometry, and the Wodzicki residue,* J. Geom. Phys., 16:327-344, 1995.
[Po] R. Ponge, *Noncommutative geometry and lower dimensional volumes in Riemannian geometry,* Lett. Math. Phys. 83 (2008), no. 1, 19–32.
[RS] S. Rempel and B. W. Schulze, Index theory of elliptic boundary problems, Akademieverlag, Berlin, 1982.
[U] W. J. Ugalde, *Differential forms and the Wodzicki residue,* J. Geom. Phys. 58 (2008), no. 12, 1739–1751.
[Wa1] Y. Wang, *Differential forms and the Wodzicki residue for manifolds with boundary,* J. Geom. Phys., 56:731-753, 2006.
[Wa2] Y. Wang, *Differential forms and the noncommutative residue for manifolds with boundary in the non-product Case,* Lett. math. Phys., 77:41-51, 2006.

[Wa3] Y. Wang, *Gravity and the noncommutative residue for manifolds with boundary,* Lett. Math. Phys. 80 (2007), no. 1, 37–56.
[Wo] M. Wodzicki, *Local invariants of spectral asymmetry*, Invent.Math. 75 no.1 143-178, 1984.
[Y] Y. Yu, *The Index Theorem and The Heat Equation Method*, Nankai Tracts in Mathematics - Vol. 2, World Scientific Publishing, 2001.

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024 , China
E-mail: *wangy581@nenu.edu.cn;*